

Regional policies targeting residential solid fuel and agricultural emissions can improve air quality and public health in the Greater Bay Area and across China

Luke Conibear^{*,1}, Carly L. Reddington¹, Ben J. Silver¹, Christoph Knote², Stephen R. Arnold¹, and Dominick V. Spracklen¹

¹ Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK

² Faculty of Medicine, University of Augsburg, Germany

* Corresponding author: Luke Conibear (L.A.Conibear@leeds.ac.uk)

Contents of this file

Supplementary Table 1: Model setup used in the Weather Research and Forecasting model online–coupled with Chemistry (WRFChem) simulations.

Supplementary Table 2: Global Exposure Mortality Model (GEMM) fit parameters for the health impact assessment from ambient fine particulate matter (PM_{2.5}) exposure (Burnett et al., 2018). All–regions, including China cohort, non–accidental function (non–communicable disease plus lower respiratory infections).

Supplementary Table 3: The impacts of policy scenarios on air quality and public health per region in China (see Supplementary Figure 2 for more details). Air quality represented by population–weighted annual–mean ambient fine particulate matter (PM_{2.5}) exposure and population–weighted maximum 6–monthly–mean daily–maximum 8–hour ambient ozone (O₃) exposure (6mDM8h). Disease burden estimates for premature mortalities (MORT, annual–sum, rounded to the nearest 100) and rate of disability–adjusted life years (DALYs, annual–mean) per 100,000 population. Scenarios are for the control (CTL) and the residential (RES), industrial within the Guangdong–Hong Kong–Macau Greater Bay Area (GBA, IND–GBA), industrial over China (IND–CHN), land transport within the GBA (TRA–GBA), land transport over China (TRA–CHN), and agriculture (AGR) scenarios relative to the control. Values in parentheses represent the 95% uncertainty intervals.

Supplementary Figure 1: Domains for Weather Research and Forecasting model online–coupled with Chemistry (WRFChem) simulations

Supplementary Figure 2: Regional groupings for (a) North China (Beijing, Tianjin, Hebei, Shanxi, and Inner Mongolia), North East China (Liaoning, Jilin, and Heilongjiang), East China (Shanghai, Jiangsu, Zhejiang, Anhui, Fujian, Jiangxi, and Shandong), South Central China (Henan, Hubei, Hunan, Guangdong, Guangxi, Hainan, Hong Kong, and Macau) including the Guangdong–Hong Kong–Macau Greater Bay Area (GBA), South West China (Chongqing, Sichuan, Guizhou, Yunnan, and Tibet), North West China (Shaanxi, Gansu, Qinghai, Ningxia, and Xinjiang), and (b) the GBA individually.

Supplementary Figure 3: Change in the annual–mean ambient fine particulate matter (PM_{2.5}) concentrations in China relative to the control (CTL) for each scenario; (a) residential (RES), (b) industry for China (IND–CHN), (c) land transport for China (TRA–CHN), and (d) agriculture (AGR).

Supplementary Figure 4: Change in the annual–mean ambient ozone (O₃) concentrations in China relative to the control (CTL) for each scenario; (a) residential (RES), (b) industry for China (IND–CHN), (c) land transport for China (TRA–CHN), and (d) agriculture (AGR).

Supplementary Figure 5: Change in the rate of disability–adjusted life years (DALYs) per 100,000 population from ambient fine particulate matter (PM_{2.5}) exposure in China relative to the control (CTL) for each scenario; (a)

residential (RES), (b) industry for China (IND-CHN), (c) land transport for China (TRA-CHN), and (d) agriculture (AGR).

Supplementary Figure 6: Change in the rate of disability-adjusted life years (DALYs) per 100,000 population from ambient ozone (O₃) exposure in China relative to the control (CTL) for each scenario; (a) residential (RES), (b) industry for China (IND-CHN), (c) land transport for China (TRA-CHN), and (d) agriculture (AGR).

Additional Supporting Information (Files uploaded separately)

The air pollution and health impact assessment data per Chinese province and GBA prefecture that support the findings of this study are available at doi.org/10.5518/919.

Supplementary Table 1: Model setup used in the Weather Research and Forecasting model online–coupled with Chemistry (WRFChem) simulations.

Model Setup and Parameterisation	
Process	Method
Timestep	180 seconds.
Horizontal	Parent grid on a resolution of 30 km along a 170×170 Lambert conformal conical grid, with a 10 km nest over Guangdong–Hong Kong–Macau Greater Bay Area (GBA).
Vertical	33 vertical levels, with 38 meteorological levels.
Microphysics	Morrison two–moment scheme (Morrison, Thompson, & Tatarskii, 2009).
Radiation	Rapid radiative transfer model for general circulation models (RRTMG), short–wave and long–wave (Iacono et al., 2008).
Boundary layer physics	Mellor–Yamada Nakanishi and Niino 2.5 (Nakanishi & Niino, 2006).
Land surface	Noah Land Surface Model (Ek et al., 2003).
Convection	Grell 3–D ensemble (Grell & Devenyi, 2002).
Gas–phase chemistry	Extended Model for Ozone and Related Chemical Tracers (MOZART, Emmons et al., 2010; A. Hodzic & Jimenez, 2011; Knote et al., 2014).
Aerosol	Updated Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) with aqueous chemistry, volatility basis set secondary organic aerosol production, and 4 sectional bins (Alma Hodzic & Knote, 2014; Knote, Hodzic, & Jimenez, 2015; Zaveri, Easter, Fast, & Peters, 2008).
Photolysis	Updated tropospheric ultraviolet–visible (TUV) photolysis based originally on Tie et al., (2003).
Dust	Global Ozone Chemistry Aerosol Radiation and Transport (GOCART) with Air Force Weather Agency (AFWA) modifications (Legrand et al., 2019).
Initial & boundary chemistry	MOZART / Goddard Earth Observing System (GEOS) Model (National Center for Atmospheric Research, 2016).
Initial & boundary meteorology	European Centre for Medium–Range Weather Forecasts (ECMWF) global reanalysis products (Dee et al., 2011).

Supplementary Table 2: Global Exposure Mortality Model (GEMM) fit parameters for the health impact assessment from ambient fine particulate matter (PM_{2.5}) exposure (Burnett et al., 2018). All-regions, including China cohort, non-accidental function (non-communicable disease plus lower respiratory infections).

Age group	θ	Standard error in θ	α	μ	ν
25+	0.1430	0.01807	1.6	15.5	36.8
25–29	0.1585	0.01477	1.6	15.5	36.8
30–35	0.1577	0.01470	1.6	15.5	36.8
35–39	0.1570	0.01463	1.6	15.5	36.8
40–44	0.1558	0.01450	1.6	15.5	36.8
45–49	0.1532	0.01425	1.6	15.5	36.8
50–54	0.1499	0.01394	1.6	15.5	36.8
55–59	0.1462	0.01361	1.6	15.5	36.8
60–64	0.1421	0.01325	1.6	15.5	36.8
65–69	0.1374	0.01284	1.6	15.5	36.8
70–74	0.1319	0.01234	1.6	15.5	36.8
75–79	0.1253	0.01174	1.6	15.5	36.8
80+	0.1141	0.01071	1.6	15.5	36.8


Supplementary Table 3: The impacts of policy scenarios on air quality and public health per region in China (see Supplementary Figure 2 for more details). Air quality represented by population-weighted annual-mean ambient fine particulate matter (PM_{2.5}) exposure and population-weighted maximum 6-monthly-mean daily-maximum 8-hour ambient ozone (O₃) exposure (6mDM8h). Disease burden estimates for premature mortalities (MORT, annual-sum, rounded to the nearest 100) and rate of disability-adjusted life years (DALYs, annual-mean) per 100,000 population. Scenarios are for the control (CTL) and the residential (RES), industrial within the Guangdong-Hong Kong-Macau Greater Bay Area (GBA, IND-GBA), industrial over China (IND-CHN), land transport within the GBA (TRA-GBA), land transport over China (TRA-CHN), and agriculture (AGR) scenarios relative to the control. Values in parentheses represent the 95% uncertainty intervals.

		CTL	RES	IND- GBA	IND- CHN	TRA- GBA	TRA- CHN	AGR
PM _{2.5} (µg m ⁻³)	China	72.8	-10.6	0.0	-0.9	0.0	-1.9	-3.2
	GBA	39.6	-1.4	+0.1	-0.5	+0.1	-0.3	-1.2
	North China	87.1	-14.3	0.0	-1.0	+0.1	-1.7	-3.8
	North East China	55.8	-10.0	+0.1	-0.6	0.0	-0.6	-1.9
	East China	74.2	-10.2	0.0	-1.1	0.0	-1.1	-3.1
	South Central China	77.6	-10.3	0.0	-0.9	0.0	-2.8	-3.7
	South West China	71.5	-11.9	+0.1	-0.7	+0.1	-2.8	-3.5
	North West China	44.7	-5.3	0.0	-0.4	0.0	-1.5	-2.1
O ₃ 6mDM8h (ppb)	China	63.5	-0.8	0.0	-0.7	0.0	+1.4	+0.5
	GBA	61.3	-0.7	0.0	-0.6	0.0	+1.0	+0.1
	North China	61.2	-0.4	0.0	-0.7	0.0	+3.2	+0.7
	North East China	54.8	-0.3	0.0	-0.5	0.0	+0.6	+0.4
	East China	60.4	-0.6	0.0	-0.9	0.0	+3.5	+0.5
	South Central China	65.7	-1.0	0.0	-0.7	0.0	+0.7	+0.4
	South West China	72.9	-1.8	0.0	-0.6	0.0	-1.6	+0.7
	North West China	63.7	-0.6	0.0	-0.3	0.0	-1.0	+0.4
PM _{2.5} MORT (yr ⁻¹)	China	2,778,700 (2,700,700 to	-188,200 (-182,900 to	+800 (+800 to	-14,300 (-13,900 to	+1,000 (+1,000 to	-33,000 (-32,100 to	-58,600 (-57,000 to


		2,864,900)	-194,000)	+900)	-14,800)	+1,000)	-34,000)	-60,400)
	GBA	104,700	-2,200	+100	-800	+100	-400	-1,900
		(101,700	(-2,100	(+100	(-800	(+100	(-400	(-1,800
		to	to	to	to	to	to	to
		107,900)	-2,300)	+100)	-800)	+100)	-400)	-1,900)
	North China	384,600	-28,900	+100	-1,700	+100	-3,400	-8,000
		(373,800	(-28,100	(+100	(-1,600	(+100	(-3,300	(-7,800
		to	to	to	to	to	to	to
		396,500)	-29,800)	+100)	-1,700)	+100)	-3,500)	-8,300)
	North East China	195,400	-21,500	+100	-1,200	+100	-1,300	-3,800
		(190,000	(-20,900	(+100	(-1,100	(+100	(-1,200	(-3,700
		to	to	to	to	to	to	to
		201,500)	-22,200)	+100)	-1,200)	+100)	-1,300)	-4,000)
	East China	851,800	-54,000	+300	-5,500	+300	-5,500	-17,200
		(827,900	(-52,500	(+300	(-5,300	(+200	(-5,400	(-16,700
		to	to	to	to	to	to	to
		878,200)	-55,700)	+300)	-5,700)	+300)	-5,700)	-17,700)
	South Central China	805,000	-44,400	+200	-3,800	+300	-12,200	-16,800
		(782,400	(-43,100	(+200	(-3,700	(+300	(-11,800	(-16,300
		to	to	to	to	to	to	to
		830,000)	-45,800)	+200)	-3,900)	+300)	-12,600)	-17,300)
	South West China	387,900	-29,400	+200	-1,500	+200	-7,700	-8,800
		(377,000	(-28,600	(+200	(-1,500	(+200	(-7,400	(-8,600
		to	to	to	to	to	to	to
		399,900)	-30,400)	+200)	-1,600)	+200)	-7,900)	-9,100)
	North West China	154,000	-10,000	0	-600	0	-2,900	-4,000
		(149,700	(-9,700	(0	(-600	(0	(-2,800	(-3,800
		to	to	to	to	to	to	to
		158,800)	-10,300)	0)	-700)	0)	-3,000)	-4,100)
PM_{2.5} DALYs rate (per 100,000 yr⁻¹)	China	4,476	-277	+1	-18	+1	-66	-99
		(3,947	(-245	(+1	(-16	(+1	(-58	(-88
		to	to	to	to	to	to	to
		5,084)	-315)	+1)	-20)	+2)	-75)	-113)
	GBA	4,887	-111	+5	-38	+5	-24	-93
		(4,309	(-98	(+4	(-33	(+5	(-21	(-82
		to	to	to	to	to	to	to
		5,551)	-126)	+5)	-43)	+6)	-28)	-106)
	North China	6,314	-469	0	-28	+2	-62	-164
		(5,569	(-414	(0	(-25	(+2	(-54	(-145
		to	to	to	to	to	to	to
		7,170)	-532)	0)	-32)	+2)	-70)	-186)
	North East China	4,900	-526	+1	-31	+1	-35	-117
		(4,321	(-464	(+1	(-27	(+1	(-31	(-103
		to	to	to	to	to	to	to
		5,566)	-598)	+1)	-35)	+1)	-40)	-133)
	East China	6,400	-391	+3	-43	+2	-49	-151
		(5,645	(-346	(+3	(-38	(+2	(-44	(-133
		to	to	to	to	to	to	to

		7,268)	−444)	+4)	−49)	+3)	−56)	−172)
	South Central China	6,358	−379	+1	−30	+2	−121	−161
		(5,608	(−335	(+1	(−26	(+2	(−107	(−142
		to	to	to	to	to	to	to
		7,221)	−430)	+2)	−34)	+3)	−138)	−183)
	South West China	5,062	−348	+2	−15	+2	−121	−126
		(4,464	(−308	(+1	(−14	(+2	(−107	(−111
		to	to	to	to	to	to	to
		5,749)	−396)	+2)	−18)	+3)	−138)	−143)
	North West China	3,883	−193	0	−13	0	−73	−97
		(3,424	(−170	(0	(−12	(0	(−64	(−86
		to	to	to	to	to	to	to
		4,411)	−219)	0)	−15)	0)	−83)	−110)
O₃ MORT (yr^{−1})	China	122,800	−3,200	0	−2,800	0	+5,800	+2,100
		(85,800	(−2,300	(0	(−2,000	(0	(+4,100	(+1,500
		to	to	to	to	to	to	to
		171,400)	−4,400)	0)	−4,400)	0)	+7,900)	+2,800)
	GBA	5,700	−200	0	−100	0	+200	0
		(4,000	(−100	(0	(−100	(0	(+100	(0
		to	to	to	to	to	to	to
		7,900)	−200)	0)	−200)	0)	+300)	0)
	North China	14,500	−200	0	−400	0	+1,700	+400
		(10,100	(−100	(0	(−300	(0	(+1,200	(+300
		to	to	to	to	to	to	to
		20,300)	−300)	0)	−500)	0)	+2,300)	+500)
	North East China	7,000	−100	0	−200	0	+200	+100
		(4,800	(−100	(0	(−100	(0	(+100	(+100
		to	to	to	to	to	to	to
		9,800)	−100)	0)	−200)	0)	+200)	+200)
	East China	32,500	−700	0	−1,100	0	+4,400	+600
		(22,700	(−500	(0	(−800	(0	(+3,100	(+400
		to	to	to	to	to	to	to
		45,500)	−1,000)	0)	−1,500)	0)	+6,000)	+800)
	South Central China	37,200	−1,100	0	−800	0	+700	+500
		(26,000	(−800	(0	(−500	(0	(+500	(+400
		to	to	to	to	to	to	to
		51,900)	−1,500)	0)	−1,000)	0)	+1,000)	+700)
	South West China	22,600	−1,000	0	−300	0	−900	+400
		(15,900	(−700	(0	(−200	(0	(−700	(+300
		to	to	to	to	to	to	to
		31,400)	−1,300)	0)	−400)	0)	−1,300)	+500)
	North West China	9,000	−200	0	−100	0	−300	+100
		(6,300	(−100	(0	(−100	(0	(−200	(+100
		to	to	to	to	to	to	to
		12,600)	−300)	0)	−100)	0)	−400)	+200)
O₃ DALYs rate	China	186	−4	0	−1	0	−6	+1
		(122	(−2	(0	(−1	(0	(−4	(+1
		to	to	to	to	to	to	to


(per 100,000 yr ⁻¹)		267)	−5)	0)	−2)	0)	−9)	+2)
	GBA	182	−5	0	−4	0	+7	0
		(120	(−3	(0	(−2	(0	(+4	(0
		to	to	to	to	to	to	to
		262)	−7)	0)	−5)	0)	+9)	+1)
	North China	184	−2	0	−4	0	+15	+4
		(120	(−1	(0	(−2	(0	(+10	(+2
		to	to	to	to	to	to	to
		263)	−3)	0)	−6)	0)	+21)	+5)
	North East China	151	−2	0	−2	0	0	+2
		(99	(−1	(0	(−1	(0	(0	(+1
		to	to	to	to	to	to	to
		218)	−2)	0)	−3)	0)	−1)	+3)
	East China	186	−3	0	−5	0	+15	+2
		(122	(−2	(0	(−3	(0	(+10	(+1
		to	to	to	to	to	to	to
		266)	−5)	0)	−7)	0)	+22)	+3)
	South Central China	216	−7	0	−4	0	−1	+2
		(142	(−5	(0	(−2	(0	(0	(+1
		to	to	to	to	to	to	to
		309)	−11)	0)	−6)	0)	−2)	+3)
	South West China	228	−9	0	−1	0	−17	+2
		(150	(−6	(0	(−1	(0	(−11	(+1
		to	to	to	to	to	to	to
		326)	−12)	0)	−2)	0)	−23)	+3)
	North West China	189	−3	0	−1	0	−10	+1
		(124	(−2	(0	(0	(0	(−6	(0
		to	to	to	to	to	to	to
		272)	−4)	0)	−1)	0)	−14)	+2)




Supplementary Figure 1: Domains for Weather Research and Forecasting model online-coupled with Chemistry (WRFChem) simulations.




Supplementary Figure 2: Regional groupings for (a) North China (Beijing, Tianjin, Hebei, Shanxi, and Inner Mongolia), North East China (Liaoning, Jilin, and Heilongjiang), East China (Shanghai, Jiangsu, Zhejiang, Anhui, Fujian, Jiangxi, and Shandong), South Central China (Henan, Hubei, Hunan, Guangdong, Guangxi, Hainan, Hong Kong, and Macau) including the Guangdong–Hong Kong–Macau Greater Bay Area (GBA), South West China (Chongqing, Sichuan, Guizhou, Yunnan, and Tibet), North West China (Shaanxi, Gansu, Qinghai, Ningxia, and Xinjiang), and (b) the GBA individually.



Supplementary Figure 3: Change in the annual-mean ambient fine particulate matter (PM_{2.5}) concentrations in China relative to the control (CTL) for each scenario; (a) residential (RES), (b) industry for China (IND-CHN), (c) land transport for China (TRA-CHN), and (d) agriculture (AGR).




Supplementary Figure 4: Change in the annual-mean ambient ozone (O_3) concentrations in China relative to the control (CTL) for each scenario; (a) residential (RES), (b) industry for China (IND-CHN), (c) land transport for China (TRA-CHN), and (d) agriculture (AGR).



Supplementary Figure 5: Change in the rate of disability-adjusted life years (DALYs) per 100,000 population from ambient fine particulate matter ($PM_{2.5}$) exposure in China relative to the control (CTL) for each scenario; (a)

residential (RES), (b) industry for China (IND-CHN), (c) land transport for China (TRA-CHN), and (d) agriculture (AGR).



Supplementary Figure 6: Change in the rate of disability-adjusted life years (DALYs) per 100,000 population from ambient ozone (O_3) exposure in China relative to the control (CTL) for each scenario; (a) residential (RES), (b) industry for China (IND-CHN), (c) land transport for China (TRA-CHN), and (d) agriculture (AGR).

References

- Burnett, R., Chen, H., Szyszkowicz, M., Fann, N., Hubbell, B., Pope, C. A., et al. (2018). Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. *Proceedings of the National Academy of Sciences*, 115(38), 9592–9597. <https://doi.org/10.1073/pnas.1803222115>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., et al. (2011). The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597. <https://doi.org/10.1002/qj.828>
- Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grunmann, P., Koren, V., et al. (2003). Implementation of Noah land surface model advances in the National Centers for Environmental Prediction operational mesoscale Eta model. *Journal of Geophysical Research: Atmospheres*, 108(D22), 8851–8867. <https://doi.org/10.1029/2002JD003296>
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., et al. (2010). Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4). *Geoscientific Model Development*, 3, 43–67. <https://doi.org/10.5194/gmd-3-43-2010>
- Grell, G. A., & Devenyi, D. (2002). A generalized approach to parameterizing convection combining ensemble and data assimilation techniques. *Geophysical Research Letters*, 29(14), 10–13. <https://doi.org/10.1029/2002GL015311>
- Hodzic, A., & Jimenez, J. L. (2011). Modeling anthropogenically controlled secondary organic aerosols in a megacity: a simplified framework for global and climate models. *Geoscientific Model Development*, 4(4), 901–917. <https://doi.org/10.5194/gmd-4-901-2011>
- Hodzic, Alma, & Knote, C. (2014). WRF-Chem 3.6.1: MOZART gas-phase chemistry with MOSAIC aerosols. *Atmospheric Chemistry Division (ACD), National Center for Atmospheric Research (NCAR)*, 7.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., & Collins, W. D. (2008). Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models. *Journal of Geophysical Research: Atmospheres*, 113(13), 2–9. <https://doi.org/10.1029/2008JD009944>
- Knote, C., Hodzic, A., & Jimenez, J. L. (2015). The effect of dry and wet deposition of condensable vapors on secondary organic aerosols concentrations over the continental US. *Atmospheric Chemistry and Physics*, 15(1), 1–18. <https://doi.org/10.5194/acp-15-1-2015>
- Knote, C., Hodzic, A., Jimenez, J. L., Volkamer, R., Orlando, J. J., Baidar, S., et al. (2014). Simulation of semi-explicit mechanisms of SOA formation from glyoxal in aerosol in a 3-D model. *Atmospheric Chemistry and Physics*, 14(12), 6213–6239. <https://doi.org/10.5194/acp-14-6213-2014>
- Legrand, S. L., Polashenski, C., Letcher, T. W., Creighton, G. A., Peckham, E., & Cetola, J. D. (2019). The AFWA Dust Emissions Scheme for the GOCART Aerosol Model in WRF-Chem. *Geoscientific Model Development*, 12, 131–166. <https://doi.org/10.5194/gmd-12-131-2019>
- Morrison, H., Thompson, G., & Tatarskii, V. (2009). Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes. *Monthly Weather Review*, 137(3), 991–1007. <https://doi.org/10.1175/2008MWR2556.1>
- Nakanishi, M., & Niino, H. (2006). An improved Mellor-Yamada Level-3 model: Its numerical stability and application to a regional prediction of advection fog. *Boundary-Layer Meteorology*, 119(2), 397–407. <https://doi.org/10.1007/s10546-005-9030-8>
- National Center for Atmospheric Research. (2016). ACOM MOZART-4/GEOS-5 global model output. UCAR. Retrieved from <http://www.acom.ucar.edu/wrf-chem/mozart.shtml>
- Tie, X., Madronich, S., Walters, S., Zhang, R., Rasch, P., & Collins, W. (2003). Effect of clouds on photolysis and oxidants in the troposphere. *Journal of Geophysical Research*, 108(D20), 4642, 1–11. <https://doi.org/10.1029/2003JD003659>
- Zaveri, R. A., Easter, R. C., Fast, J. D., & Peters, L. K. (2008). Model for Simulating Aerosol Interactions and Chemistry (MOSAIC). *Journal of Geophysical Research*, 113(D13204), 1–29. <https://doi.org/10.1029/2007JD008782>